

Computing distance-regular graph and association scheme parameters in SageMath with sage-drg

Janoš Vidali

University of Ljubljana, Ljubljana, Slovenia

janos.vidali(at)fmf.uni-lj.si

The `sage-drg` package [8] for the SageMath computer algebra system has been originally developed for computation of parameters of distance-regular graphs, and its functionality has later been extended to handle general association schemes. The package has been used to obtain nonexistence results for both distance-regular graphs and Q -polynomial association schemes, mostly using the triple intersection numbers technique, see for example [3, 4, 7].

Recently, checks for two new feasibility conditions have been implemented. The first technique, developed by Kodalen and Martin [5], relies on Schönberg's theorem on positive semidefinite functions in S^{m-1} and its application on the minimal idempotents of an association scheme. The implementation of the relevant checks in `sage-drg` allows us to replicate their nonexistence results for several feasible parameter sets for Q -polynomial association schemes.

The second technique derives from Terwilliger's work on P - and Q -polynomial association schemes [6] and has most recently been used by Gavriluk and Koolen [1, 2] to obtain some nonexistence and uniqueness results for Q -polynomial distance-regular graphs. The implementation of the relevant procedures in `sage-drg` allows us to generalize their approach and derive nonexistence for many feasible parameter sets of classical distance-regular graphs.

References

- [1] A. L. Gavriluk and J. H. Koolen, The Terwilliger polynomial of a Q -polynomial distance-regular graph and its application to pseudo-partition graphs, *Linear Algebra Appl.* **466(1)** (2015) 117–140. doi:10.1016/j.laa.2014.09.048
- [2] A. L. Gavriluk and J. H. Koolen, A characterization of the graphs of bilinear $(d \times d)$ -forms over F_2 , *Combinatorica* **39(2)** (2019) 289–321. doi:10.1007/s00493-017-3573-4
- [3] A. L. Gavriluk, S. Suda and J. Vidali, On tight 4-designs in Hamming association schemes, *Combinatorica* (2020). doi:10.1007/s00493-019-4115-z
- [4] A. L. Gavriluk, J. Vidali and J. S. Williford, On few-class Q -polynomial association schemes: feasible parameters and nonexistence results, arXiv:1908.10081 (2019).
- [5] B. G. Kodalen. Cometric Association Schemes, PhD thesis, arXiv:1905.06959 (2019).
- [6] P. Terwilliger, Lecture note on Terwilliger algebra, Edited by H. Suzuki (1993).
- [7] J. Vidali, Using symbolic computation to prove nonexistence of distance-regular graphs, *Electron. J. Combin.* **25(4):P4.21** (2018). <http://www.combinatorics.org/ojs/index.php/eljc/article/view/v25i4p21>
- [8] J. Vidali, jaanos/sage-drg: sage-drg Sage package v0.9 (2019). <https://github.com/jaanos/sage-drg/>, doi:10.5281/zenodo.3350856