



5. The 19th Century


Robin Wilson



Carl Friedrich Hindenburg (1796)




‘Hindenburg and his school [in Leipzig] attempted through systematic development of combinatorials, to give it a key position within the various mathematical disciplines.’



Peter Nicholson


(1818)

Scottish practical builder
and mathematician



Louis Poinso't's diagram-tracing puzzles (1809)

Given some points situated at random in space, it is required to arrange a single flexible thread uniting them two by two in all possible ways, so that finally the two ends of the thread join up, and so that the total length is equal to the sum of all the mutual distances. As we shall see, the solution is possible only for an odd number of points.




odd complete graphs,
 K_5, K_7, K_9, \dots ;



0123456 0246135 03625140

J. B. Listing's diagrams

Vorstudien zur Topologie (1847)



Here there are 8
odd intersections,
so we need 4 paths




This can be drawn in a single stroke, since it has only two points of odd type, both five-fold . . .



Gaston Tarry

Le problème des labyrinthes (1895)





Tarry's rule: do not return along the passage which has led to a junction for the first time unless you cannot do otherwise.

Tarry also gave a practical method for carrying this out.

W. W. Rouse Ball (1892)


Mathematical Recreations and Problems



Solving the Königsberg bridges problem corresponds to the solving the diagram-tracing puzzle on the right

In 1735 Euler did NOT draw such a picture

William Rowan Hamilton (1805-1865)




Can we visit each vertex just once and return to our starting point?

Hamilton investigates . . .

The Icosion:

A Diagram, which may also be read as a mathematical game to illustrate the Principles of the Icosian Calculus.

By Sir William Rowan Hamilton, LL.D. M. S. S. S., F. R. S. S.,
(Abstracted, from a former copy, for Section A of the British Association
Dublin, September 1st, 1857.)



$$\begin{aligned}
 1 &= i^2 = k^2 = l^2 \\
 \lambda &= i k \\
 \mu &= i k^2 \\
 \omega &= \lambda \mu \lambda \mu \\
 &= \mu \lambda \mu \lambda \mu \\
 \mu &= \lambda i \lambda \\
 \lambda &= \mu i \mu \\
 \lambda \mu^2 \lambda &= \mu \lambda \mu \\
 \mu \lambda^2 \mu &= \lambda \mu \lambda \\
 \lambda \mu^3 \lambda &= \mu^{-1} \\
 \mu \lambda^3 \mu &= \lambda^{-1} \\
 \omega^2 &= 1 \\
 \omega &= i \\
 (\lambda^3 \mu^3 (\lambda \mu)^2) &= 1.
 \end{aligned}$$

If any five consecutive points of this diagram be proposed, such as 1, 2, 3, 4, 5, or 9, 16, 15, 14, 13, it may be required to complete or finish the succession, in such a manner as to discover all the remaining fifteen points, by moving along the lines of the figure, and to end in the point close to the first of the five which were proposed. This can always be done, in at least two ways, and often in four, but never in three only, nor in more than four. Examples of such complete and cyclical successions are the following:
1, 2, 3, 4, 5, 6, 19, 18, 14, 15, 16, 17, 7, 8, 9, 10, 11, 12, 13, 20;
9, 16, 15, 14, 13, 20, 1, 2, 12, 11, 10, 3, 4, 5, 6, 19, 18, 7, 8. They have all one mathematical type, and serve as illustrations of the principles of the Icosian Calculus.
W.R.H.


icosian calculus:

$$i^2 = k^2 = l^2 = 1, \text{ where } l = ik$$

Let $m = ik^2 = lk$: then

$$\underline{l^3 m^3 l m l m l^3 m^3 l m l m = 1}$$

$l = \text{right}$
 $m = \text{left}$



Hamilton's 'quaternions':

$$i^2 = j^2 = k^2 = -1, \text{ } ijk = -1.$$


So $ij = k, ji = -k$. etc.

Hamilton's icosian game (1859)

'A Voyage Around the World'

THE ICOSIAN GAME.

Entered
at
Stationers' Hall.



Registered
agreeably to
Act V. & VI. Vic. cap. 100.

IN this new Game (invented by Sir WILLIAM ROWAN HAMILTON, LL.D., &c. of Dublin, and by him named *Icosian*, from a Greek word signifying "twenty") a player is to place the whole or part of a set of twenty numbered pieces or men upon the points or in the holes of a board, represented by the diagram above drawn in such a manner as always to proceed *along the lines* of the figure, and also to fulfil certain *other* conditions, which may in various ways be assigned by another player. Ingenuity and skill may thus be exercised in *proposing* as well as in *resolving* problems of the game. For example, the first of the two players may place the first five pieces in any five consecutive holes, and then require the second player to place the remaining fifteen men consecutively in such a manner that the succession may be *cyclical*, that is, so that No. 20 may be adjacent to No. 1; and it is always possible to answer any question of this kind. Thus, if B C D E F be the five given initial points, it is allowed to complete the succession by following the alphabetical order of the twenty consonants, as suggested by the diagram itself; but after placing the piece No. 6 in the hole H, as before, it is *also* allowed (by the supposed conditions) to put No. 7 in X instead of J, and then to conclude with the succession, W R S T V J K L M N P Q Z. Other Examples of Icosian Problems, with solutions of some of them, will be found in the following page.

LONDON:

PUBLISHED AND SOLD WHOLESALE BY JOHN JAQUES AND SON, 102 HATTON GARDEN;
AND TO BE HAD AT MOST OF THE LEADING FANCY REPOSITORIES
THROUGHOUT THE KINGDOM.


Marketing the icosian game




Hamilton sold the icosian game to a games manufacturer for £25 – a wise move, as it didn't sell.




Kirkman's 'cell of a bee' (1856)




Kirkman investigated closed paths on 'polyedra'.
But if we cut in two the cell of a bee (giving the picture above), is there such a path?



Kirchhoff's electrical networks (1845/7)



Problem: Find all the currents and voltages (using Kirchhoff's laws)




assign directions to currents

cycle VXYV: $i_1 R_1 + i_2 R_2 = E$



cycle VWYV: $i_3 R_3 + i_4 R_4 - i_2 R_2 = 0$

cycle VWYXV: $i_1 R_1 + i_3 R_3 + i_4 R_4 = E$





Fundamental set of cycles for a spanning tree

Counting trees




The six trees with 6 vertices:
how many trees have 100 vertices?




Alkanes C_nH_{2n+2} have a
tree structure: how many
have n carbon atoms?

Good Will Hunting



Draw all the homeomorphically irreducible trees with 10 vertices

Cayley's trees (1857/9)




Arthur Cayley

ON THE ANALYTICAL FORMS CALLED TREES. SECOND PART.

[From the *Philosophical Magazine*, vol. XVIII. (1859), pp. 374—378. Continuation of 203.]

THE following class of "trees" presented itself to me in some researches relating to functional symbols; viz., attending only to the terminal knots, the trees with one knot, two knots, three knots, and four knots respectively are shown in the figures 1, 2, 3 and 4:




and similarly for any number of knots. The trees with four knots are formed first from those of one knot by attaching thereto in every possible way (one way only)

Cayley's 1881 paper

To count rooted trees, Cayley used a generating function $1 + A_1x + A_2x^2 + \dots$


where $A_n =$ number of rooted trees with n branches:

he obtained the equation



$$1 + A_1x + A_2x^2 + \dots = (1-x)^{-1} \cdot (1-x^2)^{-A_1} \cdot (1-x^3)^{-A_2} \dots$$

To count unrooted trees, he started from the 'centre' of the tree and worked outwards.



n :	1	2	3	4	5	6	7	8	9	10	...
rooted:	1	2	4	9	20	48	115	286	719	1842	...
unrooted:	1	1	2	3	6	11	23	47	106	235	...

1881

On the Analytical Forms called Trees.

BY PROFESSOR CAYLEY.

In a tree of N knots, selecting any knot at pleasure as a root, the tree may be regarded as springing from this root, and it is then called a root-tree. The same tree thus presents itself in various forms as a root-tree; and if we consider the different root-trees with N knots, these are not all of them distinct trees. We have thus the two questions, to find the number of root-trees with N knots; and, to find the number of distinct trees with N knots.

I have in my paper "On the Theory of the Analytical Forms called Trees," *Phil. Mag.*, t. 13 (1857), pp. 172-176, given the solution of the first question; viz. if ϕ_N denotes the number of the root-trees with N knots, then the successive numbers $\phi_1, \phi_2, \phi_3,$ etc., are given by the formula

$$\phi_1 + x\phi_2 + x^2\phi_3 + \dots = (1-x)^{-\phi_1} (1-x^2)^{-\phi_2} (1-x^3)^{-\phi_3} \dots$$

viz. we thus find

suffix of ϕ	1	2	3	4	5	6	7	8	9	10	11	12	13
$\phi =$	1	1	2	4	9	20	48	115	286	719	1842	4766	12486

And I have, in the paper "On the Analytical Forms called Trees, with Applications to the Theory of Chemical Combinations," *Brit. Assoc. Report, 1875*, pp. 257-305, also shown how by the consideration of the centre or biceentre "of length" we can obtain formulæ for the number of central and bicentral trees; that is for the number of distinct trees, with N knots: the numerical result obtained for the total number of distinct trees with N knots is given as follows:

No. of Knots	1	2	3	4	5	6	7	8	9	10	11	12	13
No. of Central Trees	1	0	1	1	2	3	7	12	27	55	127	284	682
" Bicentral "	0	1	0	1	1	3	4	11	20	51	108	267	619
Total	1	1	1	2	3	6	11	23	47	106	235	551	1301


Counting labelled trees

1889

A THEOREM ON TREES.

[From the *Quarterly Journal of Pure and Applied Mathematics*, vol. XXIII. (1889), pp. 376—378.]

THE number of trees which can be formed with $n+1$ given knots $\alpha, \beta, \gamma, \dots$ is $=(n+1)^{n-1}$; for instance $n=3$, the number of trees with the 4 given knots $\alpha, \beta, \gamma, \delta$ is $4^2=16$, for in the first form shown in the figure the $\alpha, \beta, \gamma, \delta$ may be arranged




in 12 different orders ($\alpha\beta\gamma\delta$ being regarded as equivalent to $\delta\gamma\beta\alpha$), and in the second form any one of the 4 knots $\alpha, \beta, \gamma, \delta$ may be in the place occupied by the α : the whole number is thus $12 + 4, = 16$.

Considering for greater clearness a larger value of n , say $n=5$, I state the particular case of the theorem as follows:

No. of trees $(\alpha, \beta, \gamma, \delta, \epsilon, \zeta) =$ No. of terms of $(\alpha + \beta + \gamma + \delta + \epsilon + \zeta)^4 \alpha\beta\gamma\delta\epsilon\zeta, = 6^4, = 1296$, and it will be at once seen that the proof given for this particular case is applicable for any value whatever of n .

I use for any tree whatever the following notation: for instance, in the first of the forms shown in the figure, the branches are $\alpha\beta, \beta\gamma, \gamma\delta$; and the tree is said to be $\alpha\beta^2\gamma^2\delta$ (viz. the knots α, δ occur each once, but β, γ each twice); similarly in the second of the same forms, the branches are $\alpha\beta, \alpha\gamma, \alpha\delta$, and the tree is said




Arthur Cayley, 1889:
The number of n-vertex labelled trees is n^{n-2} .


Chemical diagrams – ‘graphic formulae’

By 1850 it was known that elements combined in fixed proportions to make compounds— formulas such as C_2H_5OH were known – but how did elements combine?

Answer: **VALENCY**



In 1864 Alexander Crum Brown introduced his ‘graphic formulae’, which then appeared in a popular textbook by Edward Frankland.



J. J. Sylvester (Nature, 1878)

... I hardly ever take up Dr. Frankland's exceedingly valuable *Notes for Chemical Students*, which are drawn up exclusively on the basis of Kekule's exquisite concept of *valence*, without deriving suggestions for new researches in the theory of algebraical forms ...

The analogy is between atoms and binary quantics exclusively. I compare every binary quantic with a chemical atom. The number of factors ... in a binary quantic is the analogue of the number of bonds, or the valence, ... of a chemical atom...

An invariant of a system of binary quantics of various degrees is the analogue of a chemical substance composed of atoms of corresponding valences... A co-variant is the analogue of an (organic or inorganic) compound radical...

radical. Every invariant and covariant thus becomes expressible by a graph precisely identical with a Kekuléan diagram or chemicograph...

Chemistry and algebra


American Journal of Mathematics, 1878



Chemistry has the same quickening and suggestive influence upon the algebraist as a visit to the Royal Academy, or the old masters may be supposed to have on a Browning or a Tennyson.

Indeed it seems to me that an exact homology exists between painting and poetry on the one hand and modern chemistry and modern algebra on the other.

Sylvester's chemical trees (1878)



J. J. Sylvester and W. K. Clifford

atom \leftrightarrow binary quantic ($ax^3 + 3bx^2y + 3cxy^2 + dy^3$)

number of bonds \leftrightarrow number of factors

chemical substance \leftrightarrow invariant (functions of a, b, c, \dots)


invariant/covariant \leftrightarrow graph


Handwritten mathematical notes illustrating the correspondence between algebraic forms and graphs:

- $\begin{vmatrix} 1 & 2 & 3 & 4 & 2 \\ 1 & 2 & 3 & 4 & 2 \end{vmatrix} =$ (graph of a cylinder)
- $4t^2 =$ (graph of a square)
- $S = \frac{1}{7} \begin{vmatrix} 1 & 2 \\ 1 & 2 \end{vmatrix} =$ (graph of a cylinder with a line)
- $R = S^2 = \frac{S}{2} \begin{vmatrix} 1 & 2 \\ 1 & 2 \end{vmatrix} =$ (graph of a cylinder with a square)
- $\begin{matrix} i \\ j \end{matrix} \begin{vmatrix} 1 & 2 & 3 & 4 & 9 \\ 1 & 2 & 3 & 4 & 6 \\ 5 & 6 & 7 & 8 & 9 \\ 5 & 6 & 7 & 8 & 6 \end{vmatrix} =$ (graph of two cylinders)
- $Cayley(2, 4, 2) =$ (graph of a square)
- $C = \square_j$
- $T = \Delta_j$
- $\frac{H}{f} \begin{vmatrix} 1 & 2 & 2 & 2 & 2 & x \\ 1 & 2 & 2 & 2 & 2 & x \end{vmatrix} =$ (graph of a cylinder with a line)
- $\frac{H}{f} \begin{vmatrix} 1 & 2 & 3 & 4 & 2 \\ 1 & 2 & 3 & 4 & 2 \end{vmatrix} =$ (graph of a triangle)
- $h = \frac{t^2}{T} \begin{vmatrix} 1 & 2 & 3 & 4 \\ 1 & 2 & 3 & 4 & 2 \end{vmatrix} =$ (graph of a cylinder with a line)
- $\{f, (i^2, A)_4\}_1$
- $\sigma = (r, f)_1$
- $1 \times x$
- $1 \times x \times x =$ (graph of a cylinder with a line)


Cayley and Sylvester's work on invariant theory was eventually supplanted by the more powerful methods of Gordan and Hilbert – the 'finite basis theorem'.

Euler diagrams (1761)







Venn diagrams (1881)







Some x are y




No x are y






All x are y



Venn diagrams for 2, 3, 4, 5 sets



Lewis Carroll's diagrams




A Syllogism worked out.

That story of yours, about your once meeting the sea-serpent, always sets me off yawning;
 I never yawn, unless when I'm listening to something totally devoid of interest.


The Premisses, separately.





The Premisses, combined.




The Conclusion.




That story of yours, about your once meeting the sea-serpent, is totally devoid of interest.





A. W. F. Edwards:
Cogwheels of the Mind



Euler's formula for plane graphs



cube



$$\begin{aligned} &(\text{number of faces}) + (\text{number of vertices}) \\ &= (\text{number of edges}) + 2 \end{aligned}$$

$$6 + 8 = 12 + 2$$


Augustin-Louis Cauchy

Möbius's problem (c.1840)




A king lay on his death-bed:
'My five sons, divide my land
among you, so that each part has
a border with each of the others.'

Möbius's problem has no
solution: five neighbouring
regions cannot exist




Dual form:

**Can you join five towns
by non-crossing roads?
no – so K_5 is non-planar**




The utilities problem (Sam Loyd, 1900)



Can we connect the three houses A, B, C to the three utilities **gas, water, electricity** without any of the connections crossing?

(Here, house B is not joined to water)

Solving the utilities problem



Is this graph $K_{3,3}$
planar?

Look at the 6-cycle
A-G-B-W-C-E-A, and try
to add the connections
A-W, G-C, and E-B . . .

